



Received on 13 January, 2014; received in revised form, 12 March, 2014; accepted, 03 May, 2014; published 01 July, 2014

FATTY ACIDS OF *HIPPOCRATEA AFRICANA* ROOT: ANTIOXIDANT AND ANTI-MICROBIAL ACTIVITY

B. N. Ita*¹ and R. Ngochindo ²

Department of Chemistry, University of Uyo ¹, Akwa Ibom State, Nigeria

Department of Pure and Industrial Chemistry ², University of Port Harcourt, Rivers State, Nigeria

Keywords:

Fatty acid content, antioxidant potential, metal chelating ability, antimicrobial activity.

Correspondence to Author:

B. N. Ita

Organic Chemistry Section,
Department of Chemistry, University
of Uyo, Akwa Ibom State, Nigeria.

E-mail: basil_ita@yahoo.com

ABSTRACT: In this study, the fatty acid content, antioxidant activities and antimicrobial potential of oil obtained from the root of *Hippocratea africana* was investigated. Results showed that the oil was characterized by oleic acid (65.73%), palmitic acid (20.34%) and linoleic acid (6.90%) as the most abundant fatty acids with a oleic acid/linoleic acid ratio of 9.52. The oil possessed potent antioxidant activity ($IC_{50} = 75.00\mu\text{g/ml}$) and exhibited significant metal chelating potentials ($EC_{50} = 7.4\text{mg/ml}$). It also showed notable antimicrobial activity against the tested human pathogens. MIC values indicated bactericidal activity against *S. aureus* (MIC =0.05mg/ml), *S. pyogenes* (MIC =0.17mg/ml) and fungicidal activity against *C. albican* (MIC =0.06mg/ml). These results support to some extent, the medicinal potential of *Hippocratea africana* root.

INTRODUCTION: *Hippocratea africana* (Willd.) Loes is a plant of the *Hippocrateaceae* family. It is a perennial climber without hairs and reproduces from seeds. The plant is widely distributed in tropical Africa. The root of the plant is used traditionally by the Ibibio's of the Niger Delta region of Nigeria in the treatment of various ailments such as fever, malaria, body pains, diabetes and diarrhoea¹. Literature survey has revealed little work on this plant species. The plant root extract has also been reported to possess *in vivo* antiplasmodial activity, analgesic, anti-inflammatory, antidiarrhoeal, antiulcer, hypolipidemic, as well as antidiabetic activity²⁻⁴.

In previous studies, isolated compounds from genus *Hippocratea* include lupeol acetate, hippocrateine I-III, mayteine, pentacyclic triterpenoids, β -sitosterol, epicatechin, friedelin, canophyllal, sesquiterpene evinonate alkaloids, as well as benzyl isothiocyanates. These compounds exhibited varying biological activities including anticancer activity, gastro protective properties, antimicrobial and anti-inflammatory activities, anti-guardial and antifeedant activities⁵⁻¹⁰. The fatty acid and antioxidant activity of tropical species of *Hippocratea africana* root has not been reported.

In this study, we report the content of fatty acids as well as the potential antioxidant and antimicrobial activity of *Hippocratea africana* root oil

MATERIALS AND METHODS:

Collection and Extraction of Oil from Test Plant
Roots of *Hippocratea africana* were collected from the wild in Uyo, Akwa Ibom State, in November, 2012, and authenticated by a taxonomist in the

	<p>QUICK RESPONSE CODE</p>
	<p>DOI: 10.13040/IJPSR.0975-8232.5(7).2732-37</p>
<p>Article can be accessed online on: www.ijpsr.com</p>	
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.5(7).2732-37</p>	

Department of Botany and Ecological Studies, University of Uyo, then dried at 40°C for two weeks. The roots were carefully derived and oil was extracted from powdered plant organ by maceration in petroleum ether (40-60°C) for 48 hours. Filtration and evaporation *in vacuo* gave yellow oil.

Determination of fatty acid profile of plant root:

The fatty acid content of the root oil was determined by gas chromatography. Fifty microlitre of oil was solubilised in 950µl of hexane and esterified using sodium ethoxide¹². The fatty acid composition was analysed by GC (Hewlett-Packard 6890) equipped with a flame ionization detector (FID), a split injector, and an HP Innowax column (30m x 0.25mm, d_f 0.25µm). The oven temperature programming was initially held at 60°C for two minutes, heated at 8°C/min up to 240°C and maintained isotherm for 15 min. The temperatures used in the injector and detector were 250 and 320°C respectively. Samples of 1µL were injected adopting a split ratio of 1:20. Hydrogen was used as the carrier gas at a linear speed of 30ml/L. Fatty acids were identified by comparison of retention times of FAME with the standard component FAME mixture.

Determination of Antioxidant potential of root oil:

The antioxidant potential of the root oil was determined by evaluating its DPPH radical scavenging activity and by measuring its metal chelating potentials.

Evaluation of DPPH Activity: Precisely 1ml of *H. africana* root oil at varying concentrations was mixed with 1ml of 0.004% methanol solution of DPPH. The mixture was shaken vigorously and allowed to stand for 30 min at room temperature in the dark. The reduction of the DPPH radical was determined by measuring the absorption at 517nm. The procedure was repeated for the blank and control. The radical scavenging activity was calculated using the equation:

DPPH scavenging effect (%)

$$= [(A_{\text{blank}} - A_{\text{sample}})/A_{\text{blank}}] \times 100.$$

Sample concentration providing EC₅₀ was calculated from the graph plotting inhibition

percentage against extract concentration¹². BHA and Vitamin E were used as positive controls.

Evaluation of Metal Chelating Activity:

Metal chelating activity was determined according to the method of Decker and Welch¹³, with some modifications. Briefly, 0.5ml of oil was mixed with 0.05ml of 2mMFeCl₂ and 0.1ml of 5mM ferrozine. The total volume was diluted with 2ml methanol. Then, the mixture was shaken vigorously and left standing at room temperature for 10mins. After the mixture had reached equilibrium, the absorbance of the solution was measured spectrophotometrically at 562nm. The percentage inhibition rate of ferrozine – Fe²⁺ complex formation was calculated using the formula:

Scavenging activity (%)

$$= [(A_{\text{control}} - A_{\text{sample}})/A_{\text{control}}] \times 100$$

Where A_{control} = absorbance of ferrozine – Fe²⁺ complex, and A_{sample} = absorbance of sample. EDTA was used as a positive control.

Antimicrobial Activity:

Antimicrobial activity of *H. africana* root oil was assayed by the disc diffusion method¹⁵. Two gram positive (*Staphylococcus aureus*, *Streptococcus pyogenes*), three gram-negative bacteria (*Salmonella typhi*, *Pseudomonas aeruginosa* and *Escherichia coli*) and a yeast (*Candida albican*) were assayed. Inocula of test isolates were prepared with fresh cultures of microbial strains and cultivated on tryptic soy agar for 24 hours at 37°C. Yeast culture was incubated on Sabouraud agar at 28°C for 48 hours.

The density of the inocula was set to 0.5McFarlands unit. Empty sterilized disks of 6mm diameter were impregnated with 1ml of oil each. 1 ml of the inocula was mixed with 17ml of Muller-Hinton agar and Sabouraud agar respectively and poured into sterile petri dishes. The disc impregnated with sample was then placed on top of the inoculated agar and pressed slightly. Plates were incubated at 37°C for 24hours (for bacterial strains) and at 28°C for 48hours (for fungal strain). Inhibition zones formed on the medium were evaluated in millimeters. Gentamicin, ciprofloxacin, penicillin and fluconazole were used as positive controls¹⁵.

Minimum inhibitory concentration (MIC) of the oil was determined using the micro-dilution broth susceptibility assay in accordance with the Clinical and Laboratory Standards Institute (NCCLS) ¹⁶. MIC is the lowest concentration of oil that inhibits the growth of microorganism.

RESULTS AND DISCUSSION: The fatty acid content of *Hippocratea africana* root oil was determined by gas chromatography. Obtained results (**Table 1**) indicate that the oil is especially characterized by oleic acid (65.73%) palmitic acid (20.34%) and linoleic acid (6.90%).

The total saturated fatty acid percentage was 25.11%, while the total amount of unsaturated fatty acid was 74.87%, of which 65.73% was monounsaturated. The oil also had an oleic acid/linoleic acid ratio of 9.52, indicating stable oil capable of preventing the formation of bad cholesterol (LDL). The high content of monounsaturated fatty acid in the plant root confers on it certain benefits. Monounsaturated fats are associated with decreased low density lipoprotein (LDL) cholesterol and possibly increased high density lipoprotein (HDL) cholesterol ¹⁶. Also,

oleic acid has been reported to control lipid oxidation through signalling/ transcriptional pathway, and may provide possibilities to treat metabolic diseases associated with lipid dysfunction ¹⁷.

It has also been reported that oleic acid decreases the content of saturated long chain fatty acids in cultured skin fibroblast from patients with adrenoleukodystrophy (ALD) ¹⁸; hence the oil may be seen as a good source of unsaturated fatty acids. Keskin and Kacar ¹⁹ reported lower values (5.78-25.7%) of oleic acid in the roots of some *Astragalus* species. *Hippocratea africana* root also contained 9.13% of EFA's (linoleic and linolenic acid). Linoleic and linolenic acids have been reported to nourish skin, hair and nails. They also help eliminate eczema, psoriasis and dandruff and help prevent hair loss ²⁰.

Our values are lower than reports for *Nyctanthes arbortristis* root ²¹ but higher than values recorded for *Vallaris solanace* (Roth) *Kuntze* root bark ²². These variations may reflect influence of factors such as vegetation, degree of ripeness, climate, soil cultivation and test methodologies.

TABLE 1: FATTY ACID PROFILE OF HIPPOCRATEA AFRICANA ROOT OIL

Fatty acid	Structure	%
Myristic	(C14:0)	0.23 ± 0.02
Palmitic	(C16:0)	20.34 ± 1.09
Stearic	(C18:0)	4.40 ± 0.21
Behenic	(C22:0)	0.03 ± 0.01
Lignoceric	(C24:0)	0.07 ± 0.01
Σ SFA		25.11 ± 1.34
Palmitoleic	(C16:1)	0.01 ± 0.01
Oleic	(C18:1)	65.73 ± 2.14
Σ MUFA		65.74 ± 2.15
Linoleic acid	(C18:2)	6.90 ± 0.62
Linolenic acid	(C18:3)	2.23 ± 0.02
Σ PUFA		9.13 ± 0.64

Antioxidant activity: DPPH assay is one of the most widely used method for evaluating the antioxidant activity of plant extracts. In this assay, the antioxidants ability to scavenge a stable radical DPPH is measured. The radical (violet colour in MeOH solution) react with suitable reducing agent, during which the electrons become paired off and the solution is progressively reduced (stoichiometrically) to a yellow coloured product,

diphenylpicrylhydrazine, with the addition of extracts in a concentration dependent manner at 517nm.

In this study, oil from the root of *Hippocratea africana* demonstrated notable antiradical activity, and this increased with increasing concentration of the oil (**Figure 1**).




FIGURE 1: DPPH SCAVENGING ACTIVITY OF HIPPOCRATEA AFRICANA ROOT OIL

The concentration at which the DPPH radical was scavenged by 50% (EC_{50}) was extrapolated and found to be 9.80mg/ml for the oil (**Table 2**). The obtained EC_{50} value for the oil was however lower than the activity of the positive controls BHA and vitamin E with EC_{50} values of 0.38 and 0.50 mg/ml respectively. The results obtained in the present work, have shown that the root oil of

TABLE 2: EC_{50} VALUES OF HIPPOCRATEA AFRICANA ROOT OIL.

	EC_{50} value* (mg/ml)	
	DPPH radical scavenging activity	Metal chelating ability
<i>Hippocratea africana</i> root oil	9.80	14.22
BHA	0.38	-
Vit. E	0.50	-
EDTA	-	0.01

* EC_{50} value, the effective concentration at which 1,1-diphenyl, 2, picrylhydrazyl (DPPH) radical were scavenged by 50%, ferrous ions were chelated by 50%, the absorbance was 0.5 for ferric reducing antioxidant power. EC_{50} value was obtained by interpolation from linear regression analysis.

Reports on the metal chelating abilities of Hippocrateaceae is rare, therefore it becomes difficult to compare our result with previous studies from the same family. However, lower chelating ability was reported for *Tabernaemontana corymbosa*²⁴, while *Ferula gummosa* Boiss roots exhibited higher chelating properties²⁵.

Antimicrobial activity: The antimicrobial activity of oil from *Hippocratea africana* root against human pathogens is shown in **Table 3**. The oil exhibited notable antimicrobial activity, and was potent against *S. aureus* (31mm, 67% inhibition), *C. albicans* (22mm, 58%inhibition) and *S. pyogenes* (16mm, 41% inhibition).

Hippocratea africana is a more effective antioxidant than the ethanolic root bark extract of *Salacia campestris* (Hippocrateaceae)²³. The observed antioxidant activity may be related to its high levels of unsaturated fatty acids and other lipophilic phytochemicals such as tocopherols, carotenoids, pigments, etc. not evaluated in this study.

Metal chelating ability: Ferrous ion is known to be good pro-oxidant and phenolic compounds have the ability to chelate pro-oxidant metal ions such as iron, thus preventing free- radical formation²⁹. The chelating ability of *H. africana* root oil increased in a dose-dependent manner (**Figure 2**). At 5mg/ml, metal chelating ability of the oil was 35.7 %; this increased to 55.6% at 20mg/ml while that of the standard tended to remain constant at higher concentration likely due to the limitation of Beer - Lamberts law. EC_{50} values (**Table 2**) indicated that the activity of the oil was lower than that of EDTA which is known to be an excellent metal chelator.




FIGURE 2: METAL CHELATING ABILITY OF HIPPOCRATEA AFRICANA ROOT OIL

An equal and more than 15mm mean zone of inhibition in the disc diffusion method were documented and the minimum inhibitory concentration (MIC) determined by the broth dilution method. MIC values indicated bactericidal activity (MIC < 0.8mg/ml) against *S. aureus*, *S. pyogenes* and *C. albicans*, but bacteriostatic (MIC = 0.8-2mg/ml) against *P. vulgaris*. Other bacterial isolates tested were resistant to the plant root. The values recorded were however less potent than the control antibiotics used. Generally, the plant exhibited better antimicrobial activity against Gram positive bacteria than Gram negative bacteria. This is in agreement with reports from Kivrak¹⁵ who

observed higher activity towards gram positive bacteria from the oils of *S. potentillifolia*. Also, Miller²⁶ reported that long chain fatty acids have higher antimicrobial activity against gram positive than gram negative bacteria. The antimicrobial activity of the plant root may in part, be attributed to its content of saturated/unsaturated fatty acids.

Lauric acid has been shown to possess antibacterial activity²⁷. Long chain unsaturated fatty acids such as linoleic and oleic acids have been reported to be bactericidal to important pathogenic microorganisms such as *S. aureus*, *H. pylori*, *C. albicans* and *E. coli*²⁸.

TABLE 3: ANTIMICROBIAL ACTIVITY OF H. AFRICANA ROOT OIL

	Oil*			Reference antibiotics*	
	Disc diffusion (mm) (%)	MIC (mg/ml)		Disc diffusion (mm) (%)	MIC (mg/ml)
<i>Staphylococcus aureus</i>	31 (67)	0.05	Penicillin	46 (100)	11
<i>Streptococcus pyogenes</i>	16 (41)	0.17	Gentamicin	39 (100)	10
<i>Salmonella typhi</i>	11 (26)	nt	Ciprofloxacin	43 (100)	8.5
<i>Escherichia coli</i>	9 (23)	nt	Penicillin	40 (100)	11.2
<i>Proteus vulgaris</i>	15 (33)	1.9	Ciprofloxacin	45 (100)	8.6
<i>Candida albicans</i>	22 (58)	0.06	Fluconazole	38 (100)	9.1

*Values are zones of inhibition in mm; (%) =percentage efficacy relative to the standard drug. nt= not tested

CONCLUSION: In the present study, the fatty acid profile, antioxidant activity and antimicrobial potential of oil obtained from the root of *Hippocratea africana* were investigated. The oil was most abundant in the unsaturated fatty acids-oleic and linoleic acid and the saturated fatty acid-palmitic acid. The oil exhibited antimicrobial activities against the tested microorganisms, but was particularly active against *Staphylococcus aureus*, *Streptococcus pyogenes* and *Candida albicans*. The plant root possessed notable antioxidant activity, which was supported by its good metal chelating abilities. Further work will focus on the fractionation and isolation of the specific compounds from the oil responsible for the observed activity.

REFERENCES:

- Okokon JE, Antia BS, Umoh EE and Etim EI: Antidiabetic and hypolipidemic effects of *Hippocratea africana* (Hippocrateaceae) in streptozotocin induced diabetic rats. *International Journal of Drug Development and Research* 2010; 2(3):501-506.
- Okokon JE, Ita BN and Udokpoh AE: The in-vivo antimalarial activities of *Uvaria chamae* and *Hippocratea africana*. *Annals of Tropical Medicine and Parasitology* 2006; 100(6):585-590.
- Okokon JE, Antia BS and Ita BN: Analgesic and anti-inflammatory effect of ethanolic root extract of *Hippocratea africana*. *International Journal of Pharmacology* 2008; 4(1):51-55.
- Okokon JE, Akpan HD, Umoh EM and Ekaidem IS: Antidiarrhoeal and antiulcer activities of *Hippocratea africana* root extract. *Pakistan Journal of Pharmaceutical Sciences*. 2011; 24(2):201-205.
- Calzada F and Mata R: Hippocrateine III: A sesquiterpene alkaloid from *Hippocratea excelsa*. *Phytochemistry* 1995; 40(2):583-585.
- Navarete A, Trejo-Miranda J and Reyes-Trejo L: Principles of root bark of *Hippocratea excelsa* (Hippocrateaceae) with gastroprotective activity. *Journal of Ethnopharmacology* 2002; 79:383-388.
- Mena-rejon GJ, Perez-Espadas AR, Moo-Puc R, Cedillo-Rivera R, Bazzocchi IL, Jimenez-Diaz IA and Quijano L: Antigiardial activity of triterpenoids from the root bark of *Hippocratea excelsa*. *Journal of Natural Products* 2007; 70(5):863-865.
- Ogbole OO, Ekor MN, Oluremi BB, Ajaiyeoba AA, Gbolade AA, Ayoola AM and Adeyemi AA: Anti-inflammatory and antimicrobial activities of *Hippocratea indica* root bark and *Poga oleosa* fruits. *African Journal of Traditional Complementary and Alternative Medicines* 2007; 4(3):372-376.

- 9 Iwu MW, Unaeze NC, Okunji CO, Corley DG, Sanson DR and Tempesta MS: Antibacterial aromatic isothiocyanates from the essential oil of *Hippocratea welwitschii* roots. *Pharmaceutical Biology* 1991; 29(2):154-158.
- 10 Christie WW: Gas chromatography and Lipids. The AOCs Lipid Library 2012; 3:44-49.
- 11 Kivrak I, Duru ME, Ozturk M, Mercan N and Harmundar GT: Antioxidants, anticholinesterase and antimicrobial constituents from the essential oil and ethanol extract of *Salvia potentillifolia*: *Food Chemistry*. 2009; 116:470-479.
- 12 Decker EA and Welch B: Role of ferritin as lipid oxidation catalyst in muscle food. *Journal of Agricultural and Food Chemistry* 1990; 38: 674-678.
- 13 Kosalec I, Kremer D, Ocatelli M, Epifano F, Genovese C, Carlucci G and Koncic, M.Z: Anthraquinone profile, antioxidant and antimicrobial activity of bark extracts of *Rhamnus alaternus*, *R. fallax*, *R. intermedia* and *R. pumila*. *Food Chemistry* 2013; 136: 335-341.
- 14 NCCLS (National Committee for Clinical Laboratory Standards). Performance standards for antimicrobial susceptibility testing: In 9th International Supplement, Wayne Pa; 1999. M100-S9.
- 15 Rudel LL, Haines JL, and Sawyer JK: Effects on plasma lipoproteins of monounsaturated, saturated and polyunsaturated fatty acids in the diet of African green monkeys. *Journal of Lipid Research* 1990; 31:1873-188
- 16 Lim JH, Gerhart-Hines Z, Dominy JE, Lee Y, Kim S and Tabata M: Oleic acid stimulates complete oxidation of fatty acids through protein kinase A-dependent activation of SIRT1-PGC1 α complex. *Journal of Biological Chemistry* 2013; 88(10):7117-7126.
- 17 Rizzo WB, Watkins PA, Phillips MW, Cranin D, Campbell B and Avigan J: Adrenoleukodystrophy: oleic acid lowers fibroblast saturated C22-26 fatty acids. *Neurology* 1986; 36(3):357-61.
- 18 Keskin C and Kacar S: Fatty acid composition of roots and shoots samples of some *Atragalus* L. (*Fabaceae*) taxa growing in the east and southeast Turkey: *Turkish Journal of Biology* 2013; 37:122-128.
- 19 Cunane S, Anderson M: Pure linoleate deficiency in the rat: Influence on growth, accumulation of n-6 polyunsaturates, and (1-¹⁴C) linoleate oxidation. *Journal of Lipid Research* 1997; 38(4):805-812.
- 20 Rahman MM and Shahjahan M: Study on fatty acid composition and physicochemical properties of *Nyctanthes arbortristis* root. *Journal of Bangladesh Chemical Society* 2011; 24(2):202-208.
- 21 Panam J, Bhanu P and Manoj G: Fatty acid composition and physico-chemical properties of oil extracted from the root of bark of *Vallan solanace* (Roth) *Kuntze*. *International Research Journal of Pharmacy* 2012; 3(12):165-166.
- 22 Velloso JCR, Khalil NM, Gutierrez VO, Macedo dos Santos VA, Furlan M and Brunetti, L: *Salacia Campestris* root bark extracts: Peroxidase inhibition, antioxidant and antiradical profiles. *Brazilian Journal of Pharmaceutical Sciences* 2009; 45(1):100-107.
- 23 Zulkefli HN, Mohamad J and Abidin NZ: Antioxidant activity of methanol extract of *Tinospora crispa* and *Tabernaemontana corymbosa*. *Sains Malaysiana* 2013; 42(6):697-706.
- 24 Ebrahimzadeh MA, Nabivi SM, Nabivi SF and Dehpour AA: Antioxidant activity of hydroalcoholic extract of *Ferula gummosa* Bioss extract. *European Review for Medical and Pharmacological Sciences* 2011; 15:658-664.
- 25 Miller RD, Brown KE and Morse SA: Inhibitory activity of fatty acids on growth of *Neisseria gonorrhoeae*. *Infections and Immunology* 1977; 17:303-306.
- 26 Ouattara B, Simard RE, Holley RA, Piette GJ and Begin A: Antibacterial activity of selected fatty acids and essential oils against six meat spoilage organisms. *International Journal of Food Microbiology* 1997; 37:155-162.
- 27 Hazell SL and Graham DY: Unsaturated fatty acids and the viability of *Helicobacter (Campylobacter) pylori*. *Journal of Clinical Microbiology* 1990; 28:1060-1061.
- 28 Bergsson G, Arnfinnsson J, Steingrimsson O and Thormar H: *In vitro* killing of *Candida albicans* by fatty acids and monoglycerides. *Antimicrobial agents and Chemotherapy* 2001; 45:3209-3212.

How to cite this article:

Ita BN and Ngochindo R: Fatty acids of *Hippocratea africana* root: antioxidant and antimicrobial activity. *Int J Pharm Sci Res* 2014; 5(7): 2732-37.doi: 10.13040/IJPSR.0975-8232.5 (7).2732-37.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)